

Datasheet for ABIN6976861

anti-NFATC3 antibody (pSer165) (AbBy Fluor® 350)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	NFATC3
Binding Specificity:	pSer165
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NFATC3 antibody is conjugated to AbBy Fluor® 350
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthesised phosphopeptide derived from human NFAT4 around the phosphorylation site of Ser165
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Cow,Sheep,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	NFATC3
---------	--------

Target Details

Alternative Name:	NFAT4 (NFATC3 Products)
Background:	Synonyms: NFAT4 (phospho S165), p-NFAT4 (phospho S165), C80703, cytoplasmic 3, D8ErtD281e, NF AT4, NF ATc3, NF-AT4, NF-ATc3, NFAC3_HUMAN, NFATc3, NFATx, Nuclear factor of activated T cells, cytoplasmic 3, nuclear factor of activated T cells, cytoplasmic, calcineurin dependent 3, Nuclear factor of activated T-cells, T cell transcription factor NFAT4, T-cell transcription factor NFAT4. Background: The product of this gene is a member of the nuclear factors of activated T cells DNA-binding transcription complex. This complex consists of at least two components: a preexisting cytosolic component that translocates to the nucleus upon T cell receptor (TCR) stimulation and an inducible nuclear component. Other members of this family participate to form this complex also. The product of this gene plays a role in the regulation of gene expression in T cells and immature thymocytes. Several transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, Nov 2010]
Gene ID:	4775
UniProt:	Q12968
Pathways:	RTK Signaling , WNT Signaling , Fc-epsilon Receptor Signaling Pathway , Chromatin Binding

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months