antibodies .- online.com







anti-C-Peptide antibody (AA 57-87) (Alexa Fluor 350)



()	1 /	0	rv	/ 1 /	71	Α.
	1//	\vdash	1 \/	16		1/1/
\sim	v	\sim	1 V	١,	_	v v

Quantity:	100 μL	
Target:	C-Peptide	
Binding Specificity:	AA 57-87	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This C-Peptide antibody is conjugated to Alexa Fluor 350	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from mouse C Peptide	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Mouse	
Purification:	Purified by Protein A.	

Target Details

Target:	C-Peptide	
Alternative Name:	C Peptide (C-Peptide Products)	

Target Details

Background:	Synonyms: proinsulin precursor, Hyperproinsulinemia, INS, Insulin Precursor, IRDN, Proinsulin,			
	Propeptide, C-Peptide, INS2_MOUSE.			
	Background: C Peptide is part of the molecule of Proinsulin, that consists of three parts: C			
	Peptide and two long strands of amino acids (called the alpha and beta chains) that later			
	become linked together to form the insulin molecule. From every molecule of proinsulin, one			
	molecule of insulin plus one molecule of C Peptide are produced. C peptide is released into the			
	blood stream in equal amounts to insulin. A test of C peptide levels will show how much insulin			
	the body is making. Insulin decreases blood glucose concentration. It increases cell			
	permeability to monosaccharides, amino acids and fatty acids. It accelerates glycolysis, the			
	pentose phosphate cycle, and glycogen synthesis in liver.			
Gene ID:	16334			
UniProt:	P01326			
Application Details				
Application Notes:	IF(IHC-P) 1:50-200			
	IF(IHC-F) 1:50-200			
	IF(ICC) 1:50-200			
Restrictions:	For Research Use only			
Handling				
Format:	Liquid			
Concentration:	1 μg/μL			
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and			
	50 % Glycerol.			
Preservative:	ProClin			
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be			
	handled by trained staff only.			
Storage:	-20 °C			
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.			
Expiry Date:	12 months			