

### Datasheet for ABIN6977096

# anti-MC1 Receptor antibody (AA 181-280) (AbBy Fluor® 350)



Go to Product page

_					
	W	0	rv	10	W

Background:

Quantity:	100 μL	
Target:	MC1 Receptor (MC1R)	
Binding Specificity:	AA 181-280	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This MC1 Receptor antibody is conjugated to AbBy Fluor® 350	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence	
	(Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human MC1 Receptor	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Purified by Protein A.	
Target Details		
Target:	MC1 Receptor (MC1R)	
Alternative Name:	MC1 Receptor (MC1R Products)	

Synonyms: MC1 Receptor, CMM5, MC1-R, MC-1R, MC1R, Melanocortin 1 receptor,

Melanocortin 1 receptor (alpha melanocyte stimulating hormone receptor), Melanocortin receptor 1, Melanocyte-stimulating hormone receptor, Melanotropin receptor, MSH-R, MSHR, MSHR\_HUMAN, SHEP2.

Background: This intronless gene encodes the receptor protein for melanocyte -stimulating hormone (MSH). The encoded protein, a seven pass transmembrane G protein coupled receptor, controls melanogenesis. Two types of melanin exist: red pheomelanin and black eumelanin. Gene mutations that lead to a loss in function are associated with increased pheomelanin production, which leads to lighter skin and hair color. Eumelanin is photoprotective but pheomelanin may contribute to UV-induced skin damage by generating free radicals upon UV radiation. Binding of MSH to its receptor activates the receptor and stimulates eumelanin synthesis. This receptor is a major determining factor in sun sensitivity and is a genetic risk factor for melanoma and non-melanoma skin cancer. Over30 variant alleles have been identified which correlate with skin and hair color, providing evidence that this gene is an important component in determining normal human pigment variation.

 Gene ID:
 4157

 UniProt:
 Q01727

Pathways: cAMP Metabolic Process, Feeding Behaviour

#### **Application Details**

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	

# Handling

Storage:	-20 °C
Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months