

#### Datasheet for ABIN6977293

# anti-SCG5 antibody (AA 1-100) (AbBy Fluor® 488)



#### Overview

| - Overview           |   |
|----------------------|---|
| Quantity:            | 100 μL  |
| Target:              | SCG5  |
| Binding Specificity: | AA 1-100  |
| Reactivity:          | Mouse, Rat  |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This SCG5 antibody is conjugated to AbBy Fluor® 488   |
| Application:         | Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |
| Product Details      |   |

| Immunogen:            | KLH conjugated synthetic peptide derived from human Secretogranin-5 |
|-----------------------|---|
| Isotype:              | IgG   |
| Cross-Reactivity:     | Mouse, Rat  |
| Predicted Reactivity: | Human,Dog,Cow,Sheep,Pig,Horse,Rabbit                                |
| Purification:         | Purified by Protein A.  |

### Target Details

| Target:           | SCG5                            |
|-------------------|---------------------------------|
| Alternative Name: | Secretogranin-5 (SCG5 Products) |

## Target Details

| 9                   |   |
|---------------------|---|
| Background:         | Synonyms: 7B2, SgV, P7B2, SGNE1, Neuroendocrine protein 7B2, Pituitary polypeptide,             |
|                     | Secretogranin V, Secretogranin-5, Secretory granule endocrine protein I, SCG5                   |
|                     | Background: Acts as a molecular chaperone for PCSK2/PC2, preventing its premature               |
|                     | activation in the regulated secretory pathway. Binds to inactive PCSK2 in the endoplasmic       |
|                     | reticulum and facilitates its transport from there to later compartments of the secretory       |
|                     | pathway where it is proteolytically matured and activated. Also required for cleavage of PCSK2  |
|                     | but does not appear to be involved in its folding. Plays a role in regulating pituitary hormone |
|                     | secretion. The C-terminal peptide inhibits PCSK2 in vitro.                                      |
| Gene ID:            | 6447  |
| UniProt:            | P05408  |
| Application Details |   |
| Application Notes:  | FCM 1:20-100  |
|                     | IF(IHC-P) 1:50-200  |
|                     | IF(IHC-F) 1:50-200  |
|                     | IF(ICC) 1:50-200  |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Format:             | Liquid  |
| Concentration:      | 1 μg/μL   |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and     |
|                     | 50 % Glycerol.  |
| Preservative:       | ProClin   |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be             |
|                     | handled by trained staff only.  |
| Storage:            | -20 °C  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.               |
| Expiry Date:        | 12 months   |
|                     |   |