antibodies .- online.com







anti-DAK antibody (AA 101-200) (Alexa Fluor 488)



Overview	
Quantity:	100 μL
Target:	DAK
Binding Specificity:	AA 101-200
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DAK antibody is conjugated to Alexa Fluor 488
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DAK
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	DAK
Alternative Name:	DAK (DAK Products)

Target Details

Background:	Synonyms: Triokinase/FMN cyclase, TKFC, Triokinase/FMN cyclase, Bifunctional ATP-
	dependent dihydroxyacetone kinase/FAD-AMP lyase (cyclizing), ATP-dependent
	dihydroxyacetone kinase, DHA kinase, Glycerone kinase, Triokinase, Triose kinase, FAD-AMP
	lyase (cyclizing), FAD-AMP lyase (cyclic FMN forming), FMN cyclase
	Background: Catalyzes both the phosphorylation of dihydroxyacetone and of glyceraldehyde,
	and the splitting of ribonucleoside diphosphate-X compounds among which FAD is the best
	substrate. Represses IFIH1-mediated cellular antiviral response.
Gene ID:	26007
UniProt:	Q3LXA3
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months