antibodies -online.com





anti-ERO1L antibody (AA 321-420) (Alexa Fluor 555)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	ER01L
Binding Specificity:	AA 321-420
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ERO1L antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Flow Cytometry (FACS)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ERO1L	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	ER01L
Alternative Name:	ERO1L (ERO1L Products)
Background: Synonyms: ER01-like protein alpha, ER01A, ER01-L, ER01-L-alpha, Endoplasmic oxidoredu	

1-like protein, Endoplasmic reticulum oxidoreductase alpha, Oxidoreductin-1-L-alpha, ERO1L Background: Oxidoreductase involved in disulfide bond formation in the endoplasmic reticulum. Efficiently reoxidizes P4HB/PDI, the enzyme catalyzing protein disulfide formation, in order to allow P4HB to sustain additional rounds of disulfide formation. Following P4HB reoxidation, passes its electrons to molecular oxygen via FAD, leading to the production of reactive oxygen species (ROS) in the cell. Required for the proper folding of immunoglobulins. Involved in the release of the unfolded cholera toxin from reduced P4HB/PDI in case of infection by V.cholerae, thereby playing a role in retrotranslocation of the toxin. Plays an important role in ER stress-induced, CHOP-dependent apoptosis by activating the inositol 1,4,5-trisphosphate receptor IP3R1.

Gene ID: 30001

UniProt: Q96HE7

Pathways: Peptide Hormone Metabolism, ER-Nucleus Signaling, Brown Fat Cell Differentiation

Application Details

Application Notes: FCM 1:20-100

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months