



[Go to Product page](#)

Datasheet for ABIN6978167

## anti-EDC4 antibody (AA 1-100) (Alexa Fluor 555)

### Overview

Quantity:	100 µL
Target:	EDC4
Binding Specificity:	AA 1-100
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EDC4 antibody is conjugated to Alexa Fluor 555
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human EDC4
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Sheep,Pig,Chicken,Rabbit
Purification:	Purified by Protein A.

### Target Details

Target:	EDC4
Alternative Name:	EDC4 ( <a href="#">EDC4 Products</a> )

## Target Details

---

Background:	<p>Synonyms: autoantigen, Autoantigen Ge 1, Autoantigen Ge-1, Autoantigen Ge1, Autoantigen RCD 8, Autoantigen RCD-8, Autoantigen RCD8, edc4, EDC4_HUMAN, Enhancer of mRNA decapping 4, Enhancer of mRNA-decapping protein 4, Ge 1, Hedls, Human enhancer of decapping large subunit, RCD 8.</p> <p>Background: The major eukaryotic mRNA decay pathway occurs through deadenylation, decapping, and 5' to 3' degradation of the mRNA. Decapping is a critical control point in this decay pathway. Edc4 (enhancer of mRNA decapping 4), also known as human enhancer of decapping large subunit (HEDLS), RCD-8 or Ge-1, is a 1,401 amino acid protein belonging to the WD repeat EDC4 family that is involved in mRNA decapping during mRNA degradation. As part of the mRNA degradation process, Edc4 becomes part of a complex that also contains hDcp1a, hDcp2a, RCK and Edc3. Localizing to P-body and cytoplasm, Edc4 contains a nuclear localization sequence (NLS) which enables it to selectively enter the nucleus as well. Edc4 becomes phosphorylated upon DNA damage and exists as two alternatively spliced isoforms that are encoded by a gene that maps to human chromosome 16q22.1.</p>
-------------	---

Gene ID:	23644
----------	-------

UniProt:	<a href="#">Q6P2E9</a>
----------	------------------------

## Application Details

---

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

---

Format:	Liquid
---------	--------

Concentration:	1 µg/µL
----------------	---------

Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
---------	---

Preservative:	ProClin
---------------	---------

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
--------------------	--

Storage:	-20 °C
----------	--------

## Handling

---

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

---

Expiry Date: 12 months