antibodies .- online.com

Datasheet for ABIN6978199 anti-VRK2 antibody (AA 151-250) (Alexa Fluor 555)



Overview

Quantity:	100 µL
Target:	VRK2
Binding Specificity:	AA 151-250
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This VRK2 antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human VRK2
lsotype:	lgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Chicken
Purification:	Purified by Protein A.
Target Details	

Target:	VRK2
Alternative Name:	VRK2 (VRK2 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN6978199 | 03/08/2024 | Copyright antibodies-online. All rights reserved.

Background:	Synonyms: Serine/threonine-protein kinase VRK2, VRK2, Vaccinia-related kinase 2
	Background: Serine/threonine kinase that regulates several signal transduction pathways.
	Isoform 1 modulates the stress response to hypoxia and cytokines, such as interleukin-1 beta
	(IL1B) and this is dependent on its interaction with MAPK8IP1, which assembles mitogen-
	activated protein kinase (MAPK) complexes. Inhibition of signal transmission mediated by the
	assembly of MAPK8IP1-MAPK complexes reduces JNK phosphorylation and JUN-dependent
	transcription. Phosphorylates 'Thr-18' of p53/TP53, histone H3, and may also phosphorylate
	MAPK8IP1. Phosphorylates BANF1 and disrupts its ability to bind DNA and reduces its binding
	to LEM domain-containing proteins. Downregulates the transactivation of transcription induced
	by ERBB2, HRAS, BRAF, and MEK1. Blocks the phosphorylation of ERK in response to ERBB2
	and HRAS. Can also phosphorylate the following substrates that are commonly used to
	establish in vitro kinase activity: casein, MBP and histone H2B, but it is not sure that this is
	physiologically relevant. Isoform 2 phosphorylates 'Thr-18' of p53/TP53, as well as histone H3.
	Reduces p53/TP53 ubiquitination by MDM2, promotes p53/TP53 acetylation by EP300 and
	thereby increases p53/TP53 stability and activity.

Gene ID:	7444
UniProt:	Q86Y07

Application Details

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN6978199 | 03/08/2024 | Copyright antibodies-online. All rights reserved.

Handling	
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months