antibodies .- online.com





anti-CILP antibody (AA 1051-1184) (Alexa Fluor 555)



| Coto | Product page |
|-------|--------------|
| G0 t0 | Froduct page |

| () | ver | VI | ew | |
|----|-----|----|----|--|
|----|-----|----|----|--|

| Quantity: | 100 μL |
|----------------------|---|
| Target: | CILP |
| Binding Specificity: | AA 1051-1184 |
| Reactivity: | Human, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This CILP antibody is conjugated to Alexa Fluor 555 |
| Application: | Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human CILP | |
|-----------------------|--|--|
| Isotype: | IgG | |
| Specificity: | The immunogen falls within the Cartilage intermediate layer protein 1 C2 region. | |
| Cross-Reactivity: | Human, Rat | |
| Predicted Reactivity: | Mouse,Cow,Sheep,Horse | |
| Purification: | Purified by Protein A. | |
| | | |

Target Details

Target: CILP

Target Details

| rarget Details | |
|---------------------|---|
| Alternative Name: | CILP (CILP Products) |
| Background: | Synonyms: CILP-1, HsT18872, Cartilage intermediate layer protein 1, Cartilage intermediate- |
| | layer protein, CILP, UNQ602/PR01188 |
| | Background: Probably plays a role in cartilage scaffolding. May act by antagonizing TGF-beta1 |
| | (TGFB1) and IGF1 functions. Has the ability to suppress IGF1-induced proliferation and sulfated |
| | proteoglycan synthesis, and inhibits ligand-induced IGF1R autophosphorylation. May inhibit |
| | TGFB1-mediated induction of cartilage matrix genes via its interaction with TGFB1. |
| | Overexpression may lead to impair chondrocyte growth and matrix repair and indirectly |
| | promote inorganic pyrophosphate (PPi) supersaturation in aging and osteoarthritis cartilage. |
| Gene ID: | 8483 |
| UniProt: | 075339 |
| Application Details | |
| Application Notes: | FCM 1:20-100 |
| | IF(IHC-P) 1:50-200 |
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
| | handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |
| | |