



[Go to Product page](#)

Datasheet for ABIN6978265

anti-MREG antibody (AA 101-200) (Alexa Fluor 555)

Overview

Quantity:	100 µL
Target:	MREG
Binding Specificity:	AA 101-200
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MREG antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DSU
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	MREG
Alternative Name:	DSU (MREG Products)

Target Details

Background: Synonyms: Dilute suppressor, Dilute suppressor protein homolog, DSU, FLJ10116, MREG_HUMAN, Melanoregulin, MGC90296, MREG, OTTHUMP00000206794, OTTHUMP00000206795, OTTHUMP00000206796, WDT 2, WDT2, Whn dependent transcript 2. Background: The photoreceptor rod cell that is responsible for vision under conditions of low light consists of stacked arrays of disk membranes that make up its outer segment portion. Regulated by complex biochemical mechanisms, the rod outer segment is under constant renewal as new disks form at the base. MREG (melanoregulin), also known as DSU (dilute suppressor protein homolog) or WDT2, is thought to play a role in membrane fusion and in regulating the biogenesis of disk membranes of photoreceptor rods. MREG interacts with RDS (also known as Peripherin-2), a photoreceptor specific tetraspanin protein that is required to maintain normal cell structure during the renewal process of membrane fusion. MREG is 214 amino acids in length, is expressed in photoreceptor cells and is expressed as two isoforms due to alternative splicing.

Gene ID: 55686

UniProt: [Q8N565](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months