
# antibodies - online.com







## anti-OSBPL3 antibody (AA 51-150) (Alexa Fluor 555)



| ( ) | ve     | K\ / |    | A . |
|-----|--------|------|----|-----|
|     | $\cup$ | 1 V/ | Щ. | V۷  |
|     |        |      |    |     |

| Quantity:            | 100 μL  |
|----------------------|---|
| Target:              | OSBPL3  |
| Binding Specificity: | AA 51-150   |
| Reactivity:          | Mouse, Rat  |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This OSBPL3 antibody is conjugated to Alexa Fluor 555   |
| Application:         | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human OSBPL3 |
|-----------------------|--|
| Isotype:              | IgG  |
| Cross-Reactivity:     | Mouse, Rat   |
| Predicted Reactivity: | Human,Dog,Cow,Horse  |
| Purification:         | Purified by Protein A.                                     |

## **Target Details**

| Target:           | OSBPL3                   |
|-------------------|--------------------------|
| Alternative Name: | OSBPL3 (OSBPL3 Products) |

#### Target Details

| Bacl | // | ro | ıın | ٨. |
|------|----|----|-----|----|
| Daci | ĸα | ΙU | uu  | u. |

Synonyms: DKFZp667P1518, KIAA0704, MGC21526, ORP 3, ORP-3, ORP3, OSBL3\_HUMAN, OSBP related protein 3, OSBP-related protein 3, OSBP3, Osbpl3, Oxysterol binding protein 3, Oxysterol binding protein like 3, Oxysterol binding protein related protein 3, Oxysterol-binding protein-related protein 3.

Background: This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has two repeats of quasi-RRM domains that bind to RNAs. It is localized in nuclear bodies of the nucleus. This protein is involved in the splicing process and it also participates in early heat shock-induced splicing arrest by transiently leaving the hnRNP complexes. Several alternatively spliced transcript variants have been noted for this gene, however, not all are fully characterized. [provided by RefSeq, Jul 2008]

Gene ID: 26031

UniProt: Q9H4L5

#### **Application Details**

Application Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

| Farmer att.        | 1:  |
|--------------------|---|
| Format:            | Liquid  |
| Concentration:     | 1 μg/μL   |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
|                    | 50 % Glycerol.  |
| Preservative:      | ProClin   |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be         |
|                    | handled by trained staff only.  |

## Handling

| Storage:         | -20 °C  |
|------------------|---|
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date:     | 12 months   |