

Datasheet for ABIN6978749

anti-GPC1 antibody (AA 251-350) (AbBy Fluor® 555)



Go to Product page

_					
	1//	r	Vİ	\triangle	۸/
	V		VI		/ V

Alternative Name:

Quantity:	100 μL	
Target:	GPC1	
Binding Specificity:	AA 251-350	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This GPC1 antibody is conjugated to AbBy Fluor® 555	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded	
	Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from mouse Glypican 1	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Mouse,Dog,Cow,Sheep,Horse	
Purification:	Purified by Protein A.	
Target Details		
Target:	GPC1	
Alta on a time Alama a	01 : 1 (00010 1 1 1)	

Glypican 1 (GPC1 Products)

Target Details

Synonyms: FLJ38078, glypican, glypican1, GPC 1, Glypican-1, GPC1, Glypican proteoglycan 1,	
Gpc1, GPC1_HUMAN, Secreted glypican-1.	
Background: Cell surface proteoglycan that bears heparan sulfate. Binds, via the heparan	
sulfate side chains, alpha-4 (V) collagen and participates in Schwann cell myelination (By	
similarity). May act as a catalyst in increasing the rate of conversion of prion protein PRPN(C)	
to PRNP(Sc) via associating (via the heparin sulfate side chains) with both forms of PRPN,	
targeting them to lipid rafts and facilitating their interaction. Required for proper skeletal muscle	
differentiation by sequestering FGF2 in lipid rafts preventing its binding to receptors (FGFRs)	
and inhibiting the FGF-mediated signaling.	
14733	
Q9QZF2	
Glycosaminoglycan Metabolic Process, Regulation of Muscle Cell Differentiation	
IF(IHC-P) 1:50-200	
IF(IHC-F) 1:50-200	
IF(ICC) 1:50-200	
For Research Use only	
Liquid	
1 μg/μL	
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
50 % Glycerol.	
ProClin	
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
handled by trained staff only.	
-20 °C	
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
12 months	