

### Datasheet for ABIN6979153

# anti-GRK2 antibody (pSer670) (AbBy Fluor® 594)



#### Overview

100 μL
GRK2 (ADRBK1)
pSer670
Human, Mouse
Rabbit
Polyclonal
This GRK2 antibody is conjugated to AbBy Fluor® 594
mmunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded
Sections) (IF (p))
KLH conjugated synthesised phosphopeptide derived from human GRK2 around the
phosphorylation site of Ser670
gG
Human, Mouse
Rat,Dog,Cow,Pig,Horse,Rabbit
Purified by Protein A.
GRK2 (ADRBK1)

## Target Details

Alternative Name:	GRK2 (ADRBK1 Products)
Background:	Synonyms: GRK2 (phospho S670), p-GRK2 (phospho S670), G-protein coupled receptor kinase
	2, ADRBK 1, ADRBK1, Adrenergic beta receptor kinase 1, BARK 1, BARK, BARK1, Beta
	adrenergic receptor kinase 1, Beta ARK 1, Beta ARK1, G Protein Coupled Receptor Kinase 2, G
	protein dependent receptor kinase 2, FLJ16718, GRK 2, ARBK1_HUMAN.
	Background: The product of this gene phosphorylates the beta-2-adrenergic receptor and
	appears to mediate agonist-specific desensitization observed at high agonist concentrations.
	This protein is an ubiquitous cytosolic enzyme that specifically phosphorylates the activated
	form of the beta-adrenergic and related G-protein-coupled receptors. Abnormal coupling of
	beta-adrenergic receptor to G protein is involved in the pathogenesis of the failing heart.
	[provided by RefSeq].
Gene ID:	156
UniProt:	P25098
Pathways:	EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Regulation of G-Protein Coupled
	Receptor Protein Signaling, CXCR4-mediated Signaling Events, G-protein mediated Events,
	Interaction of EGFR with phospholipase C-gamma, Thromboxane A2 Receptor Signaling
Application Details	
• •	JE/JUO D) 1.50 000
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

## Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months