

Datasheet for ABIN6979163

anti-KPNA4 antibody (AA 1-100) (AbBy Fluor® 594)



Go to Product page

Overview	
Quantity:	100 μL
Target:	KPNA4
Binding Specificity:	AA 1-100
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KPNA4 antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))
Product Details	
Product Details Immunogen:	KLH conjugated synthetic peptide derived from human KPNA4
	KLH conjugated synthetic peptide derived from human KPNA4
Immunogen:	
Immunogen: Isotype:	IgG
Immunogen: Isotype: Cross-Reactivity:	IgG Mouse
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity:	IgG Mouse Human,Rat,Cow,Horse,Chicken
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity: Purification:	IgG Mouse Human,Rat,Cow,Horse,Chicken

Target Details

Synonyms: Importin subunit alpha-3, KPNA4, Importin alpha Q1, Qip1, Karyopherin subunit	
alpha-4, QIP1	
Background: Functions in nuclear protein import as an adapter protein for nuclear receptor	
KPNB1. Binds specifically and directly to substrates containing either a simple or bipartite NLS	
motif. Docking of the importin/substrate complex to the nuclear pore complex (NPC) is	
mediated by KPNB1 through binding to nucleoporin FxFG repeats and the complex is	
subsequently translocated through the pore by an energy requiring, Ran-dependent	
mechanism. At the nucleoplasmic side of the NPC, Ran binds to importin-beta and the three	
components separate and importin-alpha and -beta are re-exported from the nucleus to the	
cytoplasm where GTP hydrolysis releases Ran from importin. The directionality of nuclear	
import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus. In vitro, mediates the nuclear import of human cytomegalovirus UL84 by recognizing a non-classical NLS. In vitro, mediates the nuclear	
	import of human cytomegalovirus UL84 by recognizing a nonclassical NLS.
	3840
000629	
Protein targeting to Nucleus	
IF(IHC-P) 1:50-200	
IF(IHC-F) 1:50-200	
IF(ICC) 1:50-200	
II (ICC) 1.50-200	
For Research Use only	
Liquid	
1 μg/μL	
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months