



[Go to Product page](#)

Datasheet for ABIN6979178

## anti-Importin 5 antibody (AA 501-600) (Alexa Fluor 594)

### Overview

Quantity:	100 µL
Target:	Importin 5 (IPO5)
Binding Specificity:	AA 501-600
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Importin 5 antibody is conjugated to Alexa Fluor 594
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Karyopherin beta 3
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

### Target Details

Target:	Importin 5 (IPO5)
Alternative Name:	Karyopherin beta 3 ( <a href="#">IPO5 Products</a> )

## Target Details

---

**Background:** Synonyms: DKFZp686O1576, FLJ43041, IMB3, Imp5, Importin 5, Importin beta 3 subunit, Importin subunit beta 3, Importin subunit beta-3, Importin-5, IPO 5, IPO5, IPO5\_HUMAN, Karyopherin beta 3, Karyopherin beta-3, KPNB3, MGC2068, Pse1, Ran binding protein 5, Ran GTP binding protein 5, Ran-binding protein 5, RanBP 5, RanBP5.

**Background:** Functions in nuclear protein import as nuclear transport receptor. Serves as receptor for nuclear localization signals (NLS) in cargo substrates. Is thought to mediate docking of the importin/substrate complex to the nuclear pore complex (NPC) through binding to nucleoporin and the complex is subsequently translocated through the pore by an energy requiring, Ran-dependent mechanism. At the nucleoplasmic side of the NPC, Ran binds to the importin, the importin/substrate complex dissociates and importin is re-exported from the nucleus to the cytoplasm where GTP hydrolysis releases Ran. The directionality of nuclear import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus (By similarity). Mediates the nuclear import of ribosomal proteins RPL23A, RPS7 and RPL5. Binds to a beta-like import receptor binding (BIB) domain of RPL23A. In vitro, mediates nuclear import of H2A, H2B, H3 and H4 histones. In case of HIV-1 infection, binds and mediates the nuclear import of HIV-1 Rev.

---

**Gene ID:** 3843

---

**UniProt:** [O00410](#)

---

**Pathways:** [Protein targeting to Nucleus](#)

## Application Details

---

**Application Notes:** IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

---

**Restrictions:** For Research Use only

## Handling

---

**Format:** Liquid

---

**Concentration:** 1 µg/µL

---

**Buffer:** Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

---

**Preservative:** ProClin

---

## Handling

---

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months