

## Datasheet for ABIN6979203

## anti-HS2ST1 antibody (AA 221-320) (AbBy Fluor® 594)



Go to Product page

()	ve	rvi	6	W
$\sim$	v C	1 V I	$\sim$	v v

Quantity:	100 μL
Target:	HS2ST1
Binding Specificity:	AA 221-320
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HS2ST1 antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human HS2ST1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit,Zebrafish
Purification:	Purified by Protein A.
Target Details	
Target:	HS2ST1
Alternative Name:	HS2ST1 (HS2ST1 Products)

## **Target Details**

Background:	Synonyms: 2-0-sulfotransferase, 20ST, dJ604K5.2, FLJ11317, heparan sulfate 2 0		
	sulfotransferase 1, Heparan sulfate 2-O-sulfotransferase 1, HS2ST_HUMAN, hs2st1, KIAA0448,		
	MGC131986.		
	Background: Heparan sulfate structures, which are responsible for executing multiple biologic		
	activities, are generated and regulated by heparan sulfate biosynthetic enzymes. HS2ST1 is a		
	356 amino acid single-pass type II membrane protein that localizes to the golgi apparatus and		
	belongs to the sulfotransferase 3 family. Expressed as multiple alternatively spliced isoforms,		
	HS2ST1 functions to catalyze the transfer of sulfate groups to hexuronic acid residues within		
	maturing heparan sulfate (HS), an event which is crucial for proper HS-related ligand binding		
	and signaling processes. HS2ST1 is subject to post-translational N-glycosylation and, in		
	addition to its role in HS function, may be involved in proper kidney formation.		
Gene ID:	9653		
UniProt:	Q7LGA3		
Pathways:	Glycosaminoglycan Metabolic Process, Tube Formation, SARS-CoV-2 Protein Interactome		
Application Details			
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		

	1	l:
-	ココロロ	urnen
	Hand	III IU

Expiry Date:

12 months