antibodies -online.com





anti-MED4 antibody (AA 101-200) (Alexa Fluor 594)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL
Target:	MED4
Binding Specificity:	AA 101-200
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MED4 antibody is conjugated to Alexa Fluor 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MED4
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	MED4
Alternative Name:	MED4 (MED4 Products)

Target Details

Background:	Synonyms: Mediator of RNA polymerase II transcription subunit 4, MED4, Activator-recruited
	cofactor 36 kDa component, ARC36, Mediator complex subunit 4, TRAP/SMCC/PC2 subunit
	p36 subunit, Vitamin D3 receptor-interacting protein complex 36 kDa component, DRIP36,
	MED4, ARC36, DRIP36, VDRIP, HSPC126
	Background: Component of the Mediator complex, a coactivator involved in the regulated
	transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge
	to convey information from gene-specific regulatory proteins to the basal RNA polymerase II
	transcription machinery. Mediator is recruited to promoters by direct interactions with
	regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation
	complex with RNA polymerase II and the general transcription factors.
Gene ID:	29079
UniProt:	Q9NPJ6
Pathways:	Intracellular Steroid Hormone Receptor Signaling Pathway, Nuclear Hormone Receptor Binding
	Regulation of Lipid Metabolism by PPARalpha
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

		1 .	
ш	land	linc	٩
	ıanıu	11111	J

Expiry Date:

12 months