antibodies -online.com







anti-DBNL antibody (AA 201-300) (Alexa Fluor 594)



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	DBNL	
Binding Specificity:	AA 201-300	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DBNL antibody is conjugated to Alexa Fluor 594	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HIP55
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human
Purification:	Purified by Protein A.

Target Details

Target:	DBNL
Alternative Name:	HIP55 (DBNL Products)

Target Details

Background:	Synonyms: ABP1, Actin-binding protein 1, Cervical mucin associated protein, Cervical mucin-
	associated protein, Cervical SH3P7, CMAP, Dbnl, DBNL_HUMAN, Drebrin F, drebrin like, Drebrin
	like protein, Drebrin-F, Drebrin-like protein, HIP 55, HIP-55, HPK1 interacting protein of 55 kDa,
	HPK1-interacting protein of 55 kDa, SH3 domain containing protein 7, SH3 domain-containing
	protein 7, SH3P7, Src Homology 3 Domain Containing Protein, src homology 3 domain
	containing protein HIP
	Background: Adapter protein that binds F-actin and DNM1, and thereby plays a role in receptor-
	mediated endocytosis. Plays a role in the reorganization of the actin cytoskeleton, formation of
	cell projections, such as neurites, in neuron morphogenesis and synapse formation via its
	interaction with WASL and COBL. Does not bind G-actin and promote actin polymerization by
	itself. Required for the formation of organized podosome rosettes (By similarity). May act as a
	common effector of antigen receptor-signaling pathways in leukocytes. Acts as a key
	component of the immunological synapse that regulates T-cell activation by bridging TCRs and
	the actin cytoskeleton to gene activation and endocytic processes.
Gene ID:	28988
UniProt:	Q9UJU6
Pathways:	TCR Signaling, Regulation of Actin Filament Polymerization
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months