

Datasheet for ABIN6979432

**anti-STAR antibody (AA 181-285) (AbBy Fluor® 594)**[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	STAR
Binding Specificity:	AA 181-285
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STAR antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human StAR
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Pig,Horse
Purification:	Purified by Protein A.

## Target Details

Target:	STAR
Alternative Name:	StAR ( <a href="#">STAR Products</a> )

## Target Details

---

Background:	<p>Synonyms: StARD1, Cholesterol trafficker, Luteinizing hormone induced protein, Mitochondrial steroid acute regulatory protein, StAR related lipid transfer (START) domain containing 1, StARD1, START domain containing protein 1, Steroidogenic Acute Regulatory Protein, Steroidogenic acute regulatory protein mitochondrial, STAR_HUMAN.</p> <p>Background: The protein encoded by this gene plays a key role in the acute regulation of steroid hormone synthesis by enhancing the conversion of cholesterol into pregnenolone. This protein permits the cleavage of cholesterol into pregnenolone by mediating the transport of cholesterol from the outer mitochondrial membrane to the inner mitochondrial membrane. Mutations in this gene are a cause of congenital lipid adrenal hyperplasia (CLAH), also called lipid CAH. A pseudogene of this gene is located on chromosome 13. [provided by RefSeq, Jul 2008].</p>
Gene ID:	6770
UniProt:	<a href="#">P49675</a>
Pathways:	<a href="#">Metabolism of Steroid Hormones and Vitamin D</a> , <a href="#">Response to Growth Hormone Stimulus</a> , <a href="#">C21-Steroid Hormone Metabolic Process</a> , <a href="#">Cellular Response to Molecule of Bacterial Origin</a> , <a href="#">Carbohydrate Homeostasis</a>

## Application Details

---

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

## Handling

---

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

## Handling

---

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

---

Expiry Date: 12 months