# antibodies .- online.com





## anti-DEPTOR antibody (AA 151-250) (Alexa Fluor 594)



Go to Product page

$\sim$					
	)\/e	r	٦\/	10	1///

Quantity:	100 μL	
Target:	DEPTOR	
Binding Specificity:	AA 151-250	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DEPTOR antibody is conjugated to Alexa Fluor 594	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human DEPTOR	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse	
Predicted Reactivity:	Rat,Dog,Cow,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	

### Target Details

Target:	DEPTOR
Alternative Name:	DEPTOR (DEPTOR Products)

### Target Details

Background:	Synonyms: DEP domain containing 6, DEP domain-containing mTOR-interacting protein, DEP
	domain-containing protein 6, DEPDC6, DEPTOR, DPTOR_HUMAN.
	Background: DEPTOR (DEP domain containing MTOR-interacting protein), also known as DEP.6
	or DEPDC6 (DEP domain-containing protein 6), is a 409 amino acid protein that negatively
	regulates mTORC1 and mTORC2 pathways. DEPTOR interacts with FRAP via its PDZ domain,
	and undergoes post-translational phosphorylation. Containing two DEP domains and one PDZ
	(DHR) domain, DEPTOR is encoded by a gene that maps to human chromosome 8q24.12.
	Chromosome 8 consists of nearly 146 million base pairs, encodes over 800 genes and is
	associated with a variety of diseases and malignancies. Schizophrenia, bipolar disorder,
	Trisomy 8, Pfeiffer syndrome, congenital hypothyroidism, Waardenburg syndrome and some
	leukemias and lymphomas are thought to occur as a result of defects in specific genes that
	map to chromosome 8.
Gene ID:	64798
UniProt:	Q8TB45
Pathways:	PI3K-Akt Signaling, RTK Signaling, Regulation of Cell Size, BCR Signaling, Warburg Effect
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C

## Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months