

[Go to Product page](#)

Datasheet for ABIN6979805

anti-CD101 antibody (AA 51-150) (Alexa Fluor 647)

Overview

| | |
|----------------------|---|
| Quantity: | 100 µL |
| Target: | CD101 |
| Binding Specificity: | AA 51-150 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This CD101 antibody is conjugated to Alexa Fluor 647 |
| Application: | Flow Cytometry (FACS), Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|---|
| Immunogen: | KLH conjugated synthetic peptide derived from human CD101 |
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Predicted Reactivity: | Mouse,Rat |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|--|
| Target: | CD101 |
| Alternative Name: | CD101 (CD101 Products) |

Target Details

| | |
|-------------|--|
| Background: | <p>Synonyms: V7, IGSF2, EWI-101, Immunoglobulin superfamily member 2, Cell surface glycoprotein V7, Glu-Trp-Ile EWI motif-containing protein 101, CD101, EWI101</p> <p>Background: Plays a role as inhibitor of T-cells proliferation induced by CD3. Inhibits expression of IL2RA on activated T-cells and secretion of IL2. Inhibits tyrosine kinases that are required for IL2 production and cellular proliferation. Inhibits phospholipase C-gamma-1/PLCG1 phosphorylation and subsequent CD3-induced changes in intracellular free calcium. Prevents nuclear translocation of nuclear factor of activated T-cell to the nucleus. Plays a role in the inhibition of T-cell proliferation via IL10 secretion by cutaneous dendritic cells. May be a marker of CD4(+) CD56(+) leukemic tumor cells.</p> |
|-------------|--|

| | |
|----------|------|
| Gene ID: | 9398 |
|----------|------|

| | |
|----------|------------------------|
| UniProt: | Q93033 |
|----------|------------------------|

Application Details

| | |
|--------------------|--|
| Application Notes: | FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 |
|--------------------|--|

| | |
|---------------|-----------------------|
| Restrictions: | For Research Use only |
|---------------|-----------------------|

Handling

| | |
|--------------------|--|
| Format: | Liquid |
| Concentration: | 1 µg/µL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |