antibodies .- online.com





anti-CYP8B1 antibody (AA 151-250) (AbBy Fluor® 647)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL
Target:	CYP8B1
Binding Specificity:	AA 151-250
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CYP8B1 antibody is conjugated to AbBy Fluor® 647
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CYP8B1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Cow,Sheep,Horse
Purification:	Purified by Protein A.

Target Details

Target:	CYP8B1
Alternative Name:	CYP8B1 (CYP8B1 Products)

Target Details

Precaution of Use:

Storage Comment:

Storage:

Expiry Date:

Target Details		
Background:	Synonyms: 7-alpha-hydroxy-4-cholesten-3-one 12-alpha-hydroxylase, 7-alpha-hydroxycholest-4-	
	en-3-one 12-alpha-hydroxylase, CP8B1_HUMAN, CYP12, CYP8B1, CYPVIIIB1, Cytochrome P450	
	8B1, Cytochrome P450, family 8, subfamily B, polypeptide 1, Sterol 12-alpha-hydroxylase.	
	Background: This gene encodes a member of the cytochrome P450 superfamily of enzymes.	
	The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved	
	in drug metabolism and synthesis of cholesterol, steroids and other lipids. This endoplasmic	
	reticulum membrane protein catalyzes the conversion of 7 alpha-hydroxy-4-cholesten-3-one	
	into 7-alpha,12-alpha-dihydroxy-4-cholesten-3-one. The balance between these two steroids	
	determines the relative amounts of cholic acid and chenodeoxycholic acid both of which are	
	secreted in the bile and affect the solubility of cholesterol. This gene is unique among the	
	cytochrome P450 genes in that it is intronless. [provided by RefSeq, Jul 2008]	
Gene ID:	1582	
UniProt:	Q9UNU6	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

handled by trained staff only.

-20 °C

12 months

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be