antibodies .- online.com





anti-MLZE antibody (AA 1-100) (Alexa Fluor 647)



Go to Product page

| \sim | | | |
|--------|-----|-----|-----|
| | N/P | r\/ | i⊢₩ |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | MLZE | |
| Binding Specificity: | AA 1-100 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This MLZE antibody is conjugated to Alexa Fluor 647 | |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human GSDMC |
|-------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | Purified by Protein A. |

Target Details

| Target: | MLZE |
|-------------------|--|
| Alternative Name: | GSDMC (MLZE Products) |
| Background: | Synonyms: Gasdermin C, Gasdermin-C, GSDMC, GSDMC_HUMAN, Melanoma derived leucine |

Target Details

Expiry Date:

12 months

| l arget Details | |
|---------------------|---|
| | zipper, extra nuclear factor, Melanoma-derived leucine zipper-containing extranuclear factor, |
| | MLZE |
| | Background: The N-terminal moiety promotes pyroptosis. May be acting by homooligomerizing |
| | within the membrane and forming pores (PubMed:27281216). The physiological relevance of |
| | this observation is unknown (Probable). |
| Gene ID: | 56169 |
| UniProt: | Q9BYG8 |
| Application Details | |
| Application Notes: | IF(IHC-P) 1:50-200 |
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
| | handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |