

[Go to Product page](#)

Datasheet for ABIN6979988

**anti-TMEM30A antibody (AA 251-350) (Alexa Fluor 647)**

## Overview

|                      |  |
|----------------------|--|
| Quantity:            | 100 µL   |
| Target:              | TMEM30A  |
| Binding Specificity: | AA 251-350   |
| Reactivity:          | Human, Mouse, Rat  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This TMEM30A antibody is conjugated to Alexa Fluor 647   |
| Application:         | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Western Blotting (WB) |

## Product Details

|                       |   |
|-----------------------|---|
| Immunogen:            | KLH conjugated synthetic peptide derived from human Transmembrane protein 30A |
| Isotype:              | IgG   |
| Cross-Reactivity:     | Human, Mouse, Rat   |
| Predicted Reactivity: | Cow,Pig,Rabbit  |
| Purification:         | Purified by Protein A.  |

## Target Details

|                   |  |
|-------------------|--|
| Target:           | TMEM30A  |
| Alternative Name: | Transmembrane protein 30A ( <a href="#">TMEM30A Products</a> ) |

## Target Details

|             |  |
|-------------|--|
| Background: | <p>Synonyms: CDC50A, C6orf67, Cell cycle control protein 50A, P4-ATPase flippase complex beta subunit TMEM30A, Transmembrane protein 30A, TMEM30A</p> <p>Background: Accessory component of a P4-ATPase flippase complex which catalyzes the hydrolysis of ATP coupled to the transport of aminophospholipids from the outer to the inner leaflet of various membranes and ensures the maintenance of asymmetric distribution of phospholipids. Phospholipid translocation seems also to be implicated in vesicle formation and in uptake of lipid signaling molecules. The beta subunit may assist in binding of the phospholipid substrate. Required for the proper folding, assembly and ER to Golgi exit of the ATP8A2:TMEM30A flippase complex. ATP8A2:TMEM30A may be involved in regulation of neurite outgrowth, and, reconstituted to liposomes, predominantly transports phosphatidylserine (PS) and to a lesser extent phosphatidylethanolamine (PE). The ATP8A1:TMEM30A flippase complex seems to play a role in regulation of cell migration probably involving flippase-mediated translocation of phosphatidylethanolamine (PE) at the plasma membrane. Required for the formation of the ATP8A2, ATP8B1 and ATP8B2 P-type ATPase intermediate phosphoenzymes. Involved in uptake of platelet-activating factor (PAF), synthetic drug alkylphospholipid edelfosine, and, probably in association with ATP8B1, of perifosine. Also mediate the export of alpha subunits ATP8A1, ATP8B1, ATP8B2, ATP8B4, ATP10A, ATP10B, ATP10D, ATP11A, ATP11B and ATP11C from the ER to other membrane localizations.</p> |
|-------------|--|

|          |       |
|----------|-------|
| Gene ID: | 55754 |
|----------|-------|

|          |                        |
|----------|------------------------|
| UniProt: | <a href="#">Q9NV96</a> |
|----------|------------------------|

## Application Details

|                    |  |
|--------------------|--|
| Application Notes: | IF(IHC-P) 1:50-200<br>IF(IHC-F) 1:50-200<br>IF(ICC) 1:50-200 |
|--------------------|--|

|               |                       |
|---------------|-----------------------|
| Restrictions: | For Research Use only |
|---------------|-----------------------|

## Handling

|         |        |
|---------|--------|
| Format: | Liquid |
|---------|--------|

|                |         |
|----------------|---------|
| Concentration: | 1 µg/µL |
|----------------|---------|

|         |   |
|---------|---|
| Buffer: | Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
|---------|---|

## Handling

---

|                    |  |
|--------------------|--|
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |