

#### Datasheet for ABIN6980056

# anti-HIPK1 antibody (AA 1-100) (AbBy Fluor® 647)



Go to Product page

_					
	W	0	rv	10	W

Overview		
Quantity:	100 μL	
Target:	HIPK1	
Binding Specificity:	AA 1-100	
Reactivity:	Human, Mouse, Monkey, Macaque	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This HIPK1 antibody is conjugated to AbBy Fluor® 647	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human HIPK1	
Isotype:	IgG	
Cross-Reactivity:	Human, Macague, Monkey, Mouse	

Cross-Reactivity: Human, Macaque, Monkey, Mouse

Predicted Reactivity: Rat, Dog, Cow, Sheep, Pig, Horse, Rabbit

Purification: Purified by Protein A.

### **Target Details**

Target: HIPK1

Alternative Name: HIPK1 (HIPK1 Products)

### Target Details

Background:	Synonyms: Hipk 1, Hipk1, HIPK1_HUMAN, Homeodomain interacting protein kinase 1,	
	Homeodomain interacting protein kinase 1 like protein, Homeodomain-interacting protein	
	kinase 1, Myak, Nbak 2, Nbak2, Nuclear body associated kinase 2, Nuclear body associated	
	kinase 2b	
	Background: Serine/threonine-protein kinase involved in transcription regulation and TNF-	
	mediated cellular apoptosis. Plays a role as a corepressor for homeodomain transcription	
	factors. Phosphorylates DAXX and MYB. Phosphorylates DAXX in response to stress, and	
	mediates its translocation from the nucleus to the cytoplasm. Inactivates MYB transcription	
	factor activity by phosphorylation. Prevents MAP3K5-JNK activation in the absence of TNF.	
	TNF triggers its translocation to the cytoplasm in response to stress stimuli, thus activating	
	nuclear MAP3K5-JNK by derepression and promoting apoptosis. May be involved in anti-	
	oxidative stress responses. Involved in the regulation of eye size, lens formation and retinal	
	lamination during late embryogenesis. Promotes angiogenesis and to be involved in erythroid	
	differentiation. May be involved in malignant squamous cell tumor formation.	
Gene ID:	204851	
UniProt:	Q86Z02	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	

## Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months