antibodies -online.com





anti-NEK6 antibody (AA 81-180) (AbBy Fluor® 647)



Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL
Target:	NEK6
Binding Specificity:	AA 81-180
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NEK6 antibody is conjugated to AbBy Fluor® 647
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NEK6
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	NEK6
Alternative Name:	NEK6 (NEK6 Products)

Target Details

Storage Comment:

Background:	Synonyms: Highly similar to cell cycle protein kinase CDC5/MSD2, NEK6, NEK6_HUMAN, Never
	in mitosis A-related kinase 6, NIMA (Never In Mitosis Gene A) Related Kinase 6, NimA related
	protein kinase 6, NimA-related protein kinase 6, Protein kinase SID6 1512, Protein kinase SID6-
	1512, Putative serine threonine protein kinase, Serine/threonine protein kinase Nek6,
	Serine/threonine-protein kinase Nek6, SID61512.
	Background: Protein kinase which plays an important role in mitotic cell cycle progression.
	Required for chromosome segregation at metaphase-anaphase transition, robust mitotic
	spindle formation and cytokinesis. Phosphorylates ATF4, CIR1, PTN, RAD26L, RBBP6, RPS7,
	RPS6KB1, TRIP4, STAT3 and histones H1 and H3. Phosphorylates KIF11 to promote mitotic
	spindle formation. Involved in G2/M phase cell cycle arrest induced by DNA damage. Inhibition
	of activity results in apoptosis. May contribute to tumorigenesis by suppressing p53/TP53-
	induced cancer cell senescence.
Gene ID:	10783
UniProt:	Q9HC98
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

		1.	
\vdash	land	lın	
	ıaııu	1111	U

Expiry Date:

12 months