# antibodies -online.com





## anti-ORC2 antibody (AA 201-300) (Alexa Fluor 680)



Go to Product page

| $\sim$ |    |    |    |             |   |
|--------|----|----|----|-------------|---|
| ()     | VE | ۲۱ | /1 | $\triangle$ | Λ |

| Quantity:            | 100 μL  |  |
|----------------------|---|--|
| Target:              | ORC2  |  |
| Binding Specificity: | AA 201-300  |  |
| Reactivity:          | Mouse   |  |
| Host:                | Rabbit  |  |
| Clonality:           | Polyclonal  |  |
| Conjugate:           | This ORC2 antibody is conjugated to Alexa Fluor 680   |  |
| Application:         | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |  |

## Product Details

| Immunogen:            | KLH conjugated synthetic peptide derived from human ORC2 |
|-----------------------|--|
| Isotype:              | IgG  |
| Cross-Reactivity:     | Mouse  |
| Predicted Reactivity: | Human,Rat,Dog,Cow,Sheep,Horse,Chicken,Rabbit             |
| Purification:         | Purified by Protein A.                                   |

## **Target Details**

| Target:           | ORC2                 |
|-------------------|----------------------|
| Alternative Name: | ORC2 (ORC2 Products) |

#### **Target Details**

Optional[synonyms]: The origin recognition complex (ORC) is a highly conserved six subunits protein complex essential for the initiation of the DNA replication in eukaryotic cells. Studies in yeast demonstrated that ORC binds specifically to origins of replication and serves as a platform for the assembly of additional initiation factors such as Cdc6 and Mcm proteins. The protein encoded by this gene is a subunit of the ORC complex. This protein forms a core complex with ORC3, -4, and -5. It also interacts with CDC45 and MCM10, which are proteins known to be important for the initiation of DNA replication. This protein has been demonstrated to specifically associate with the origin of replication of Epstein-Barr virus in human cells, and is thought to be required for DNA replication from viral origin of replication. Alternatively spliced transcript variants have been found, one of which is a nonsense-mediated mRNA decay candidate. [provided by RefSeq, Oct 2010]

Gene ID:

4999

UniProt:

Q13416

### **Application Details**

Application Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

| Format:            | Liquid   |  |
|--------------------|--|--|
| Concentration:     | 1 μg/μL  |  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |  |
| Preservative:      | ProClin  |  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |  |
| Storage:           | -20 °C   |  |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |  |
| Expiry Date:       | 12 months  |  |