# antibodies -online.com





# anti-NAPRT1 antibody (AA 351-450) (AbBy Fluor® 680)



Go to Product page

( )	11/0	K\ /	iew	1
	$\cup$	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	NAPRT1 (NAPRT)	
Binding Specificity:	AA 351-450	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This NAPRT1 antibody is conjugated to AbBy Fluor® 680	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human NAPRT1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

## **Target Details**

Target:	NAPRT1 (NAPRT)
Alternative Name:	NAPRT1 (NAPRT Products)

## **Target Details**

•		
Background:	Synonyms: FHA HIT interacting protein, FHA-HIT-interacting protein, FHIP, FHIP, NAPRT1,	
	NAPRTase, Nicotinate phosphoribosyltransferase, Nicotinate phosphoribosyltransferase	
	domain containing 1, Nicotinate phosphoribosyltransferase domain-containing protein 1,	
	Nicotinic acid phosphoribosyltransferase, PNCB_HUMAN, PP3856.	
	Background: Nicotinic acid (NA, niacin) is converted by nicotinic acid	
	phosphoribosyltransferase (NAPRT, EC 2.4.2.11) to NA mononucleotide (NaMN), which is then	
	converted to NA adenine dinucleotide (NaAD), and finally to nicotinamide adenine dinucleotide	
	(NAD), which serves as a coenzyme in cellular redox reactions and is an essential component	
	of a variety of processes in cellular metabolism including response to stress (Hara et al.,	
	2007).[supplied by OMIM, Mar 2008]	
Gene ID:	93100	
UniProt:	Q6XQN6	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	