antibodies -online.com





anti-DAK antibody (AA 101-200) (Alexa Fluor 680)



()	1 /	\sim	KI /	110	Νę
	1//	\vdash	I \/	1 ←	٠// ٢

Quantity:	100 μL	
Target:	DAK	
Binding Specificity:	AA 101-200	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DAK antibody is conjugated to Alexa Fluor 680	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DAK	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	DAK
Alternative Name:	DAK (DAK Products)

Target Details

3		
Background:	Synonyms: Triokinase/FMN cyclase, TKFC, Triokinase/FMN cyclase, Bifunctional ATP-	
	dependent dihydroxyacetone kinase/FAD-AMP lyase (cyclizing), ATP-dependent	
	dihydroxyacetone kinase, DHA kinase, Glycerone kinase, Triokinase, Triose kinase, FAD-AMP	
	lyase (cyclizing), FAD-AMP lyase (cyclic FMN forming), FMN cyclase	
	Background: Catalyzes both the phosphorylation of dihydroxyacetone and of glyceraldehyde,	
	and the splitting of ribonucleoside diphosphate-X compounds among which FAD is the best	
	substrate. Represses IFIH1-mediated cellular antiviral response.	
Gene ID:	26007	
UniProt:	Q3LXA3	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	