antibodies .- online.com



Datasheet for ABIN6981272

anti-Solute Carrier Family 35 (UDP-GlcNAc/UDP-Glucose Transporter), Member D2 (SLC35D2) (AA 251-326) antibody (Alexa Fluor 680)



Go to Product page

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL	
Target:	Solute Carrier Family 35 (UDP-GlcNAc/UDP-Glucose Transporter), Member D2 (SLC35D2)	
Binding Specificity:	AA 251-326	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	Alexa Fluor 680	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from mouse SLC35D2
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Pig
Purification:	Purified by Protein A.

Target Details

Target:	Solute Carrier Family 35 (UDP-GlcNAc/UDP-Glucose Transporter), Member D2 (SLC35D2)	
Alternative Name:	SLC35D2 (SLC35D2 Products)	

Target Details

_		
Background:	Synonyms: UDP-N-acetylglucosamine/UDP-glucose/GDP-mannose transporter, Slc35d2, Solute	
	carrier family 35 member D2, UDP-galactose transporter-related protein 8, UGTrel8, Slc35d2,	
	Ugtrel8	
	Background: Antiporter transporting nucleotide sugars such as UDP-N-acetylglucosamine	
	(UDP-GlcNAc), UDP-glucose (UDP-Glc) and GDP-mannose (GDP-Man) pooled in the cytosol into	
	the lumen of the Golgi in exchange for the corresponding nucleosides monophosphates (UMP	
	for UDP-sugars and GMP for GDP-sugars). May take part in heparan sulfate synthesis by	
	supplying UDP-Glc-NAc, the donor substrate, and thus be involved in growth factor signaling.	
Gene ID:	70484	
UniProt:	Q762D5	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	