

[Go to Product page](#)

Datasheet for ABIN6981789

**anti-DTX3L antibody (AA 651-740) (Alexa Fluor 750)**

## Overview

Quantity:	100 µL
Target:	DTX3L
Binding Specificity:	AA 651-740
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DTX3L antibody is conjugated to Alexa Fluor 750
Application:	Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DTX3L
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	DTX3L
Alternative Name:	DTX3L ( <a href="#">DTX3L Products</a> )

## Target Details

Background:	<p>Synonyms: BBAP, E3 ubiquitin-protein ligase DTX3L, B-lymphoma- and BAL-associated protein, Protein deltex-3-like, Rhysin-2, Rhysin2, DTX3L</p> <p>Background: Ubiquitin ligase that mediates monoubiquitination of 'Lys-91' of histone H4 (H4K91ub1), in response to DNA damage. Protects cells exposed to DNA-damaging agents. The exact role of H4K91ub1 in DNA damage response is still unclear but it may function as a licensing signal for additional histone H4 post-translational modifications such as H4 'Lys-20' methylation (H4K20me). Involved in the recruitment of 53BP1/TP53BP1 to sites of DNA damage by mediating H4K91ub1 formation. In concert with PARP9, plays a role in PARP1-dependent DNA damage repair. PARP1-dependent PARP9-DTX3L-mediated ubiquitination promotes the rapid and specific recruitment of 53BP1/TP53BP1, UIMC1/RAP80, and BRCA1 to DNA damage sites.</p>
Gene ID:	151636
UniProt:	<a href="#">Q8TDB6</a>
Pathways:	<a href="#">Notch Signaling</a>

## Application Details

Application Notes:	FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

Handling

---

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months