antibodies -online.com





anti-PTPN21 antibody (AA 171-270) (Alexa Fluor 750)



Go to Product page

| \sim | | | |
|--------|-----|-----|-----|
| | N/P | r\/ | i⊢₩ |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | PTPN21 |
| Binding Specificity: | AA 171-270 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This PTPN21 antibody is conjugated to Alexa Fluor 750 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human PTPD1 | |
|-----------------------|---|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human, Mouse, Rat | |
| Predicted Reactivity: | Dog,Cow,Pig,Horse,Rabbit | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | PTPN21 |
|-------------------|-------------------------|
| Alternative Name: | PTPD1 (PTPN21 Products) |

Target Details

| Pookground: | Cynonyma: Drotain tyrogina phagabataga D1. Drotain tyrogina phagabataga nan ragartar tyrogina | | | |
|---------------------|--|--|--|--|
| Background: | Synonyms: Protein tyrosine phosphatase D1, Protein tyrosine phosphatase non receptor type | | | |
| | 21, PTPN 21, PTPN21, PTN21_HUMAN, PTPNL, PTPRL10, Tyrosine protein phosphatase non | | | |
| | receptor type 21. | | | |
| | Background: The protein encoded by this gene is a member of the protein tyrosine phosphatase | | | |
| | (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP contains an N-terminal domain, similar to cytoskeletal- associated proteins including band 4.1, ezrin, merlin, and radixin. This PTP was shown to specially interact with BMX/ETK, a member of Tec tyrosine kinase family characterized by a multimodular structures including PH, | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | SH3, and SH2 domains. The interaction of this PTP with BMX kinase was found to increase the | | | |
| | activation of STAT3, but not STAT2 kinase. Studies of the similar gene in mice suggested the | | | |
| | possible roles of this PTP in liver regeneration and spermatogenesis. [provided by RefSeq, Jul | | | |
| | 2008] | | | |
| Gene ID: | 11099 | | | |
| UniProt: | Q16825 | | | |
| Application Details | | | | |
| | IE(IIIO D) 1.F0 200 | | | |
| Application Notes: | IF(IHC-P) 1:50-200 | | | |
| | IF(IHC-F) 1:50-200 | | | |
| | IF(ICC) 1:50-200 | | | |
| Restrictions: | For Research Use only | | | |
| Handling | | | | |
| Format: | Liquid | | | |
| Concentration: | 1 μg/μL | | | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. | | | |
| Preservative: | ProClin | | | |
| - 10301 VALIVE. | | | | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. | | | |
| Storage: | -20 °C | | | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | | | |
| | | | | |

| | | 1. | |
|----------|-------|------|---|
| \vdash | land | lın | |
| | ıaııu | 1111 | U |
| | | | |

Expiry Date:

12 months