

Datasheet for ABIN6982667

anti-NFKBIA antibody (pSer32) (Biotin)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	NFKBIA
Binding Specificity:	pSer32
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NFKBIA antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human phospho-IKB alpha (Ser32)
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Cow, Sheep, Pig, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	NFKBIA
Alternative Name:	IKB alpha (NFKBIA Products)

Target Details

Background:	Synonyms: IKB alpha (phospho S32), IKB alpha (phospho Ser32), p-IKB alpha (phospho S32), I kappa B alpha, I-kappa-B-alpha, IkappaBalpna, Ikb-alpha, IKBA, IKBA_HUMAN, IKBalpha, MAD 3, MAD3, Major histocompatibility complex enhancer-binding protein MAD3, NF kappa B inhibitor alpha, NF-kappa-B inhibitor alpha, NFKBI, NFKBIA, Nuclear factor of kappa light chain gene enhancer in B cells, Nuclear factor of kappa light polypeptide gene enhancer in B cells inhibitor alpha. Background: This gene encodes a member of the NF-kappa-B inhibitor family, which contain multiple ankrin repeat domains. The encoded protein interacts with REL dimers to inhibit NF-kappa-B/REL complexes which are involved in inflammatory responses. The encoded protein moves between the cytoplasm and the nucleus via a nuclear localization signal and CRM1-mediated nuclear export. Mutations in this gene have been found in ectodermal dysplasia anhidrotic with T-cell immunodeficiency autosomal dominant disease. [provided by RefSeq, Aug 2011]
Gene ID:	4792
UniProt:	P25963
Pathways:	NF-kappaB Signaling , TCR Signaling , TLR Signaling , Fc-epsilon Receptor Signaling Pathway , Activation of Innate immune Response , Cellular Response to Molecule of Bacterial Origin , Maintenance of Protein Location , Hepatitis C , Protein targeting to Nucleus , Toll-Like Receptors Cascades , BCR Signaling

Application Details

Application Notes:	IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months