

[Go to Product page](#)

Datasheet for ABIN6982679 **anti-LSM6 antibody (Biotin)**

Overview

Quantity:	100 µL
Target:	LSM6
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LSM6 antibody is conjugated to Biotin
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human LSM6:31-80/80
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Purification:	Purified by Protein A.

Target Details

Target:	LSM6
Alternative Name:	LSM6 (LSM6 Products)
Background:	<p>Synonyms: LSM 6, LSM6 homolog U6 small nuclear RNA associated, LSM6 homolog, U6 small nuclear RNA associated (S. cerevisiae), Sm protein F, U6 snRNA associated Sm like protein LSM6, YDR378C.</p> <p>Background: Sm-like proteins were identified in a variety of organisms based on sequence</p>

Target Details

homology with the Sm protein family (see SNRPD2, MIM 601061). Sm-like proteins contain the Sm sequence motif, which consists of 2 regions separated by a linker of variable length that folds as a loop. The Sm-like proteins are thought to form a stable heteromer present in tri-snRNP particles, which are important for pre-mRNA splicing.[supplied by OMIM, Apr 2004]

Function : Component of LSm protein complexes, which are involved in RNA processing and may function in a chaperone-like manner, facilitating the efficient association of RNA processing factors with their substrates. Component of the cytoplasmic LSM1-LSM7 complex, which is thought to be involved in mRNA degradation by activating the decapping step in the 5'-to-3' mRNA decay pathway. Component of the nuclear LSM2-LSM8 complex, which is involved in splicing of nuclear mRNAs. LSM2-LSM8 associates with multiple snRNP complexes containing the U6 snRNA (U4/U6 di-snRNP, spliceosomal U4/U6.U5 tri-snRNP, and free U6 snRNP). It binds directly to the 3'-terminal U-tract of U6 snRNA and plays a role in the biogenesis and stability of the U6 snRNP and U4/U6 snRNP complexes. LSM2-LSM8 probably also is involved degradation of nuclear pre-mRNA by targeting them for decapping, and in processing of pre-tRNAs, pre-rRNAs and U3 snoRNA. Subunit : Component of the heptameric LSM1-LSM7 complex, which consists of LSM1, LSM2, LSM3, LSM4, LSM5, LSM6 and LSM7. Component of the heptameric LSM2-LSM8 complex, which consists of LSM2, LSM3, LSM4, LSM5, LSM6, LSM7 and LSM8. The LSm subunits form a seven-membered ring structure with a doughnut shape. Similarity : Belongs to the snRNP Sm proteins family. SmF/LSm6 subfamily.

Gene ID: 11157

UniProt: [P62312](#)

Application Details

Application Notes: IHC-P 1:200-400

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months