

[Go to Product page](#)

Datasheet for ABIN6982727

**anti-SMARCC1 antibody (AA 301-400) (Biotin)**

## Overview

Quantity:	100 µL
Target:	SMARCC1
Binding Specificity:	AA 301-400
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SMARCC1 antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SMARCC1
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human, Mouse, Dog, Cow, Sheep, Horse, Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	SMARCC1
Alternative Name:	SMARCC1 ( <a href="#">SMARCC1 Products</a> )

## Target Details

Background:	<p>Synonyms: AI115498, BAF 155, BAF155, BRG 1 associated factor 155, BRG1 associated factor 155, BRG1-associated factor 155, Chromatin remodeling complex BAF155 subunit, CRACC 1, CRACC1, Mammalian chromatin remodeling complex BRG 1 associated factor 155, Mammalian chromatin remodeling complex BRG1 associated factor 155, Rsc 8, Rsc8, SMARCC1, SMARCC 1, SMARCC1, SMRC1_HUMAN, SRG 3, SRG3, SWI 3, SWI/SNF complex 155 kDa subunit, SWI/SNF complex subunit SMARCC1, SWI/SNF related matrix associated actin dependent regulator of chromatin c1, SWI/SNF related matrix associated actin dependent regulator of chromatin subfamily c member 1, SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily C member 1, SWI3.</p> <p>Background: Involved in transcriptional activation and repression of select genes by chromatin remodeling (alteration of DNA-nucleosome topology). May stimulate the ATPase activity of the catalytic subunit of the complex. Also involved in vitamin D-coupled transcription regulation via its association with the WINAC complex, a chromatin-remodeling complex recruited by vitamin D receptor (VDR), which is required for the ligand-bound VDR-mediated transrepression of the CYP27B1 gene. Belongs to the neural progenitors-specific chromatin remodeling complex (npBAF complex) and the neuron-specific chromatin remodeling complex (nBAF complex). During neural development a switch from a stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The transition from proliferating neural stem/progenitor cells to post-mitotic neurons requires a switch in subunit composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes (nBAF). The npBAF complex is essential for the self-renewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along with CREST plays a role regulating the activity of genes essential for dendrite growth.</p>
Gene ID:	601732
UniProt:	<a href="#">Q92922</a>
Pathways:	<a href="#">Chromatin Binding</a>

## Application Details

Application Notes:	IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months