# antibodies -online.com





## anti-CILP antibody (AA 1051-1184) (Cy3)



Go to Product page

_			
$\cup$	)ve	rv	iew

Quantity:	100 μL
Target:	CILP
Binding Specificity:	AA 1051-1184
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CILP antibody is conjugated to Cy3
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human CILP	
Isotype:	IgG	
Specificity:	The immunogen falls within the Cartilage intermediate layer protein 1 C2 region.	
Cross-Reactivity:	Human, Rat	
Predicted Reactivity:	Mouse,Cow,Sheep,Horse	
Purification:	Purified by Protein A.	

#### **Target Details**

Target:	CILP

### **Target Details**

rarger betaile		
Alternative Name:	CILP (CILP Products)	
Background:	Synonyms: CILP-1, HsT18872, Cartilage intermediate layer protein 1, Cartilage intermediate-	
	layer protein, CILP, UNQ602/PR01188	
	Background: Probably plays a role in cartilage scaffolding. May act by antagonizing TGF-beta1	
	(TGFB1) and IGF1 functions. Has the ability to suppress IGF1-induced proliferation and sulfated	
	proteoglycan synthesis, and inhibits ligand-induced IGF1R autophosphorylation. May inhibit	
	TGFB1-mediated induction of cartilage matrix genes via its interaction with TGFB1.	
	Overexpression may lead to impair chondrocyte growth and matrix repair and indirectly	
	promote inorganic pyrophosphate (PPi) supersaturation in aging and osteoarthritis cartilage.	
Gene ID:	8483	
UniProt:	075339	
Application Details		
Application Notes:	FCM 1:20-100	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	