antibodies -online.com





anti-MOXD1 antibody (AA 501-600) (Cy5)



Go to Product page

| () | 11/0 | K\ / | iew |
|-----|--------|------|-------------|
| | \cup | ועוי | $I \cap VV$ |
| | | | |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | MOXD1 | |
| Binding Specificity: | AA 501-600 | |
| Reactivity: | Human, Mouse | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This MOXD1 antibody is conjugated to Cy5 | |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human MOXD1 | |
|-----------------------|---|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human, Mouse | |
| Predicted Reactivity: | Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | MOXD1 |
|-------------------|------------------------|
| Alternative Name: | MOXD1 (MOXD1 Products) |

Target Details

| Background: | Synonyms: DBH like monooxygenase protein 1, DBH-like monooxygenase protein 1, dJ248E1.1, | | |
|---------------------|--|--|--|
| | DKFZp564G202, Monooxygenase X, monooxygenase, DBH like 1, MOX, Moxd1, | | |
| | MOXD1_HUMAN, PRO5780. | | |
| | Background: MOXD1 is a 613 amino acid single-pass type I membrane protein of the | | |
| | Endoplasmic reticulum that belongs to the copper type II ascorbate-dependent | | |
| | monooxygenase family. Existing as two alternatively spliced isoforms, MOXD1 is expressed in | | |
| | adult spinal cord, adrenal gland, brain, testis, uterus, lung and kidney, as well as fetal liver and | | |
| | brain. MOXD1 is upregulated during replicative senescence in primary fibroblast and umbilical | | |
| | vein endothelial cell cultures, and uses two copper ions per subunit as a cofactor. MOXD1 | | |
| | contains one DOMON domain, undergoes post-translational N-glycosylation and is encoded by | | |
| | a gene that maps to human chromosome 6. Chromosome 6 contains 170 million base pairs, | | |
| | comprises nearly 6 % of the human genome and is associated with early onset intestinal | | |
| | cancer, Porphyria cutanea tarda, Parkinson's disease and Stickler syndrome. | | |
| Gene ID: | 26002 | | |
| UniProt: | Q6UVY6 | | |
| Application Details | | | |
| | | | |
| Application Notes: | IF(IHC-P) 1:50-200 | | |
| | IF(IHC-F) 1:50-200 | | |
| | IF(ICC) 1:50-200 | | |
| Restrictions: | For Research Use only | | |
| Handling | | | |
| Format: | Liquid | | |
| Concentration: | 1 μg/μL | | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | | |
| | 50 % Glycerol. | | |
| Preservative: | ProClin | | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | | |
| | handled by trained staff only. | | |
| Storage: | -20 °C | | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | | |
| | | | |

| \vdash | land | lına |
|----------|-------|------|
| | iaria | шц |

Expiry Date:

12 months