# antibodies -online.com





## anti-CILP antibody (AA 1051-1184) (Cy5.5)



Go to Product page

/ N	1	-	K	/1	01	A /
0	1//	$\vdash$	1 \	/ I	-	<b>\/\/</b>
$\sim$	٧.	$\sim$	1 V		$\sim$	v v

Quantity:	100 μL
Target:	CILP
Binding Specificity:	AA 1051-1184
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CILP antibody is conjugated to Cy5.5
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	nmunogen: KLH conjugated synthetic peptide derived from human CILP	
Isotype:	IgG	
Specificity:	The immunogen falls within the Cartilage intermediate layer protein 1 C2 region.	
Cross-Reactivity:	Human, Rat	
Predicted Reactivity:	Mouse,Cow,Sheep,Horse	
Purification:	Purified by Protein A.	

#### **Target Details**

Target:	CILP	

### **Target Details**

rarger betano	
Alternative Name:	CILP (CILP Products)
Background:	Synonyms: CILP-1, HsT18872, Cartilage intermediate layer protein 1, Cartilage intermediate-
	layer protein, CILP, UNQ602/PRO1188
	Background: Probably plays a role in cartilage scaffolding. May act by antagonizing TGF-beta1
	(TGFB1) and IGF1 functions. Has the ability to suppress IGF1-induced proliferation and sulfated
	proteoglycan synthesis, and inhibits ligand-induced IGF1R autophosphorylation. May inhibit
	TGFB1-mediated induction of cartilage matrix genes via its interaction with TGFB1.
	Overexpression may lead to impair chondrocyte growth and matrix repair and indirectly
	promote inorganic pyrophosphate (PPi) supersaturation in aging and osteoarthritis cartilage.
Gene ID:	8483
UniProt:	075339
Application Details	
Application Notes:	FCM 1:20-100
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months