

Datasheet for ABIN6985137

anti-GRK2 antibody (pSer670) (Cy5.5)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	GRK2 (ADRBK1)
Binding Specificity:	pSer670
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GRK2 antibody is conjugated to Cy5.5
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthesised phosphopeptide derived from human GRK2 around the phosphorylation site of Ser670
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	GRK2 (ADRBK1)
---------	---------------

Target Details

Alternative Name:	GRK2 (ADRBK1 Products)
Background:	Synonyms: GRK2 (phospho S670), p-GRK2 (phospho S670), G-protein coupled receptor kinase 2, ADRBK 1, ADRBK1, Adrenergic beta receptor kinase 1, BARK 1, BARK, BARK1, Beta adrenergic receptor kinase 1, Beta ARK 1, Beta ARK1, G Protein Coupled Receptor Kinase 2, G protein dependent receptor kinase 2, FLJ16718, GRK 2, ARBK1_HUMAN. Background: The product of this gene phosphorylates the beta-2-adrenergic receptor and appears to mediate agonist-specific desensitization observed at high agonist concentrations. This protein is an ubiquitous cytosolic enzyme that specifically phosphorylates the activated form of the beta-adrenergic and related G-protein-coupled receptors. Abnormal coupling of beta-adrenergic receptor to G protein is involved in the pathogenesis of the failing heart. [provided by RefSeq].
Gene ID:	156
UniProt:	P25098
Pathways:	EGFR Signaling Pathway , Neurotrophin Signaling Pathway , Regulation of G-Protein Coupled Receptor Protein Signaling , CXCR4-mediated Signaling Events , G-protein mediated Events , Interaction of EGFR with phospholipase C-gamma , Thromboxane A2 Receptor Signaling

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months