

Datasheet for ABIN6985726

anti-STAT3 antibody (pTyr705) (Cy5.5)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	STAT3
Binding Specificity:	pTyr705
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This STAT3 antibody is conjugated to Cy5.5
Application:	Immunofluorescence (Cultured Cells) (IF (cc))

Product Details

Immunogen:	KLH conjugated Synthesised phosphopeptide derived from human STAT3 around the phosphorylation site of Tyr705
Clone:	3F10
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified by Protein G.

Target Details

Target:	STAT3
Alternative Name:	STAT3 (STAT3 Products)

Target Details

Background: Synonyms: STAT3 (phospho Y705), p-STAT3 (phospho Y705), Phospho-Stat3_pTyr705? STAT3(Phospho-Tyr705), p-STAT3(Tyr705), Phosphorylated Stat3(pTyr705), p-Stat3, Acute Phase Response Factor, APRF, DNA binding protein APRF, FLJ20882, MGC16063, Signal Transducer and Activator of Transcription 3, STAT 3, STAT3_HUMAN.

Background: The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. This protein mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of this protein. PIAS3 protein is a specific inhibitor of this protein. Three alternatively spliced transcript variants encoding distinct isoforms have been described. Gene Symbol: APRF, FLJ20882, HIES, MGC16063.

Gene ID:	6774
UniProt:	P40763
Pathways:	JAK-STAT Signaling , RTK Signaling , Interferon-gamma Pathway , Neurotrophin Signaling Pathway , Dopaminergic Neurogenesis , Response to Growth Hormone Stimulus , Carbohydrate Homeostasis , Stem Cell Maintenance , Hepatitis C , Protein targeting to Nucleus , Feeding Behaviour , CXCR4-mediated Signaling Events , Signaling of Hepatocyte Growth Factor Receptor

Application Details

Application Notes:	IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months