antibodies -online.com





anti-CCDC64 antibody (AA 401-500) (FITC)



Go to Product page

| \sim | | | |
|--------|-----|-----|-----|
| | N/P | r\/ | i⊢₩ |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | CCDC64 |
| Binding Specificity: | AA 401-500 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This CCDC64 antibody is conjugated to FITC |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human CCDC64 |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Predicted Reactivity: | Mouse,Rat,Cow,Pig,Horse,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | CCDC64 |
|-------------------|--------------------------|
| Alternative Name: | CCDC64 (CCDC64 Products) |

Target Details

| 5 | | |
|---------------------|--|--|
| Background: | Synonyms: BICD family-like cargo adapter 1, BICDL1, Bicaudal D-related protein 1, BICD-related | |
| | protein 1, BICDR-1, Coiled-coil domain-containing protein 64A, CCDC64A, BICDR1, CCDC64 | |
| | Background: Component of secretory vesicle machinery in developing neurons that acts as a | |
| | regulator of neurite outgrowth. Regulates the secretory vesicle transport by controlling the | |
| | accumulation of Rab6-containing secretory vesicles in the pericentrosomal region restricting | |
| | anterograde secretory transport during the early phase of neuronal differentiation, thereby | |
| | inhibiting neuritogenesis. | |
| Gene ID: | 92558 | |
| UniProt: | Q6ZP65 | |
| Application Details | | |
| Application Notes: | IF(IHC-P) 1:50-200 | |
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | |
| | 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| | handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |