

## Datasheet for ABIN6986805

## anti-ERO1LB antibody (AA 1-100) (FITC)



$\sim$					
	W	0	rv	10	W

Alternative Name:

Quantity:	100 μL	
Target:	ERO1LB	
Binding Specificity:	AA 1-100	
Reactivity:	Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ERO1LB antibody is conjugated to FITC	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human ERO1LB	
Isotype:	IgG	
Cross-Reactivity:	Mouse, Rat	
Predicted Reactivity:	Human,Pig,Rabbit,Guinea Pig	
Purification:	Purified by Protein A.	
Target Details		
Target:	ERO1LB	

**ERO1LB (ERO1LB Products)** 

## **Target Details**

Background:	Synonyms: Endoplasmic oxidoreductin-1-like protein B, ER01-L-beta, ER01B, ER01-like protein			
	beta, Oxidoreductin-1-L-beta			
	Background: Oxidoreductase involved in disulfide bond formation in the endoplasmic reticulum.			
	Efficiently reoxidizes P4HB/PDI, the enzyme catalyzing protein disulfide formation, in order to			
	allow P4HB to sustain additional rounds of disulfide formation. Other protein disulfide			
	isomerase family members can also be reoxidized, but at lower rates compared to P4HB,			
	including PDIA2 (50 % of P4HB reoxidation rate), as well as PDIA3, PDIA4, PDIA6 and NXNDC12			
	(<10 %). Following P4HB reoxidation, passes its electrons to molecular oxygen via FAD, leading			
	to the production of reactive oxygen species (ROS) in the cell. May be involved in oxidative			
	proinsulin folding in pancreatic cells, hence may play a role in glucose homeostasis.			
Cana ID:				
Gene ID:	56605			
UniProt:	Q86YB8			
Pathways:	SARS-CoV-2 Protein Interactome			
Application Details				
Application Notes:	IF(IHC-P) 1:50-200			
	IF(IHC-F) 1:50-200			
	IF(ICC) 1:50-200			
Restrictions:	For Research Use only			
Handling				
Format:	Liquid			
Concentration:	1 μg/μL			
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and			
	50 % Glycerol.			
Preservative:	ProClin			
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be			
	handled by trained staff only.			
Storage:	-20 °C			
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.			
Expiry Date:	12 months			