antibodies -online.com





anti-FIBIN antibody (AA 101-200) (FITC)



Overview

Quantity:	100 μL
Target:	FIBIN
Binding Specificity:	AA 101-200
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FIBIN antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FIBIN
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	FIBIN
Alternative Name:	FIBIN (FIBIN Products)

Target Details

Expiry Date:

Background:	Synonyms: Fin bud initiation factor homolog, FIBIN, PSEC0235
	Background: FIBIN (Fin bud initiation factor homolog) is a 211 amino acid protein involved in fin
	initiation in zebrafish. The human homolog is encoded by a gene that maps to chromosome 11,
	which makes up around 4 $\%$ of human genomic DNA and is considered a gene and disease
	association dense chromosome. The chromosome 11 encoded Atm gene is important for
	regulation of cell cycle arrest and apoptosis following double strand DNA breaks. Atm mutation
	leads to the disorder known as ataxia-telangiectasia. The blood disorders Sickle cell anemia
	and $_$ thalassemia are caused by HBB gene mutations. Wilms' tumors, WAGR syndrome and
	Denys-Drash syndrome are associated with mutations of the WT1 gene. Jervell and Lange-
	Nielsen syndrome, Jacobsen syndrome, Niemann-Pick disease, hereditary angioedema and
	Smith-Lemli-Opitz syndrome are also associated with defects in chromosome 11.
Gene ID:	387758
UniProt:	Q8TAL6
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

12 months