

[Go to Product page](#)

Datasheet for ABIN6986900
anti-SBNO1 antibody (AA 331-430) (FITC)

Overview

Quantity:	100 µL
Target:	SBNO1
Binding Specificity:	AA 331-430
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SBNO1 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SBNO1
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit,Zebrafish
Purification:	Purified by Protein A.

Target Details

Target:	SBNO1
Alternative Name:	SBNO1 (SBNO1 Products)

Target Details

Background:	<p>Synonyms: FLJ10701, FLJ10833, FLJ16176, Monocyte protein 3, MOP 3, MOP-3, MOP3, Protein strawberry notch homolog 1, SBNO 1, Sbnol, SBNO1_HUMAN, Sno, Sno strawberry notch homolog 1, Strawberry notch homolog 1.</p> <p>Background: SBNO1 is a 1,392 amino acid protein encoded by the human gene of the same name located on chromosome 12. Encoding over 1,100 genes within 132 million bases, chromosome 12 makes up about 4.5 % of the human genome. A number of skeletal deformities are linked to chromosome 12, including hypochondrogenesis, achondrogenesis and Kniest dysplasia. Noonan syndrome, which includes heart and facial developmental defects among the primary symptoms, is caused by a mutant form of PTPN11 gene product, SH-PTP2. Chromosome 12 is also home to a homeobox gene cluster, which encodes crucial transcription factors for morphogenesis, and the natural killer complex gene cluster, encoding C-type lectin proteins which mediate the NK cell response to MHC I interaction. Trisomy 12p leads to facial development defects, seizure disorders and a host of other symptoms which vary in severity depending on the extent of mosaicism. It is most severe in cases of complete trisomy.</p>
Gene ID:	55206
UniProt:	A3KN83
Pathways:	SARS-CoV-2 Protein Interactome

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months