

Datasheet for ABIN6987016

anti-MED12 antibody (AA 251-350) (FITC)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	MED12
Binding Specificity:	AA 251-350
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MED12 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MED12/Trap230
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Cow,Sheep,Pig,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	MED12
Alternative Name:	MED12/Trap230 (MED12 Products)

Target Details

Background:	Synonyms: Mediator of RNA polymerase II transcription subunit 12, MED12, Activator-recruited cofactor 240 kDa component, ARC240, CAG repeat protein 45 Mediator complex subunit 12, OPA-containing protein, Thyroid hormone receptor-associated protein complex 230 kDa component, Trap230, Trinucleotide repeat-containing gene 11 protein, MED12, ARC240, CAGH45, HOPA, KIAA0192, TNRC11, TRAP230 Background: Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. This subunit may specifically regulate transcription of targets of the Wnt signaling pathway and SHH signaling pathway.
Gene ID:	9968
UniProt:	Q93074
Pathways:	Intracellular Steroid Hormone Receptor Signaling Pathway , Nuclear Hormone Receptor Binding , Stem Cell Maintenance , Chromatin Binding , Regulation of Lipid Metabolism by PPARalpha , Tube Formation

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months