antibodies - online.com





anti-UGCG antibody (AA 21-120) (FITC)



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	UGCG	
Binding Specificity:	AA 21-120	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This UGCG antibody is conjugated to FITC	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Ceramide glucosyltransferase
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	UGCG
Alternative Name:	Ceramide glucosyltransferase (UGCG Products)

Target Details

Synonyms: CEGT_HUMAN, UGCG, GCS, GLCT 1, GLCT-1, GLCT1, Glucosylceramide synthase,
UDP glucose ceramide glucosyltransferase, UDP glucose N acylsphingosine D
glucosyltransferase, UDP-glucose ceramide glucosyltransferase, UDP-glucose:N-
acylsphingosine D-glucosyltransferase, Ugcg.
Background: Catalyzes the first glycosylation step in glycosphingolipid biosynthesis, the
transfer of glucose to ceramide. May also serve as a "flippase".
7357
Q16739
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
For Research Use only
Liquid
1 μg/μL
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
50 % Glycerol.
ProClin
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
handled by trained staff only.
-20 °C
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
12 months