

Datasheet for ABIN6987297

anti-Caspase 12 antibody (AA 21-100) (FITC)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	Caspase 12 (CASP12)
Binding Specificity:	AA 21-100
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Caspase 12 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Caspase-12
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human
Purification:	Purified by Protein A.

Target Details

Target:	Caspase 12 (CASP12)
Alternative Name:	Caspase-12 (CASP12 Products)

Target Details

Background:	Synonyms: Caspase-12, UNQ9415, Apoptosis related cysteine protease, CASP 12, casp12, CASP12P1, Caspase 12 pseudogene 1, CASP-12, casp12, caspase 12 (gene/pseudogene), Caspase12, CASPC_HUMAN, OTTHUMP00000207026, OTTHUMP00000207028, OTTHUMP00000207031, OTTHUMP00000207032, OTTHUMP00000219058, UNQ9415. Background: Caspases are cysteine proteases that cleave C-terminal aspartic acid residues on their substrate molecules. This gene is most highly related to members of the ICE subfamily of caspases that process inflammatory cytokines. In rodents, the homolog of this gene mediates apoptosis in response to endoplasmic reticulum stress. However, in humans this gene contains a polymorphism for the presence or absence of a premature stop codon. The majority of human individuals have the premature stop codon and produce a truncated non-functional protein. The read-through codon occurs primarily in individuals of African descent and carriers have endotoxin hypo-responsiveness and an increased susceptibility to severe sepsis. Several alternatively spliced transcript variants have been noted for this gene. [provided by RefSeq, Feb 2011]
Gene ID:	100506742
UniProt:	Q6UXS9
Pathways:	Apoptosis , ER-Nucleus Signaling , Positive Regulation of Endopeptidase Activity , Unfolded Protein Response

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months