# antibodies .- online.com





# anti-STAG2 antibody (AA 1021-1120) (HRP)



Go to Product page

( )	11/0	K\ /	iew	1
	$\cup$	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	STAG2
Binding Specificity:	AA 1021-1120
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STAG2 antibody is conjugated to HRP
Application:	Western Blotting (WB)

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human SA2
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

# Target Details

Target:	STAG2
Alternative Name:	SA2 (STAG2 Products)
Background:	Synonyms: Cohesin Subunit SA 2, Cohesin subunit SA-2, DKFZp686P168, DKFZp781H1753, SA

2, SA2, SCC3 homolog 2, STAG 2, STAG 2, stag2, STAG2\_HUMAN, Stromal antigen 2, FLJ25871, bA51701.1.

Background: Component of cohesin complex, a complex required for the cohesion of sister chromatids after DNA replication. The cohesin complex apparently forms a large proteinaceous ring within which sister chromatids can be trapped. At anaphase, the complex is cleaved and dissociates from chromatin, allowing sister chromatids to segregate. The cohesin complex may also play a role in spindle pole assembly during mitosis.

Gene ID: 10735

UniProt: Q8N3U4

Pathways: Stem Cell Maintenance

### **Application Details**

Application Notes: WB 1:300-5000

Restrictions: For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months