

Datasheet for ABIN6988517

anti-Caspase 9 antibody (AA 231-330) (PE)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	Caspase 9 (CASP9)
Binding Specificity:	AA 231-330
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Caspase 9 antibody is conjugated to PE
Application:	Western Blotting (WB), Flow Cytometry (FACS)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Caspase-9
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Cow, Sheep, Pig, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	Caspase 9 (CASP9)
Alternative Name:	Caspase-9 (CASP9 Products)
Background:	Synonyms: Caspase-9 subunit p35, Apaf-3, APAF 3, APAF3, Apoptosis related cysteine

Target Details

peptidase, Apoptotic protease activating factor 3, Apoptotic protease MCH 6, Apoptotic protease MCH6, CASP 9, CASP9, Caspase 9, Caspase 9 apoptosis related cysteine protease, Caspase 9 precursor, Caspase 9c, Caspase9, Caspase9 subunit p10, ICE LAP6, ICE like apoptotic protease 6, RNCASP9, MCH 6, MCH6, OTTHUMP00000044594, CASP9_HUMAN. Background: Caspase 9 (also known as ICE like apoptotic protease 6 (ICE LAP6), apoptotic protease Mch6, and apoptotic protease activating factor 3 (Apaf3)) is a member of the peptidase family C14 that contains a CARD domain. This caspase is active as a heterotetramer and has been reported to have two isoforms. ProCaspase 9 has been reported to be approximately 47 kD. This caspase is present in the cytosol and, upon activation, translocates to the mitochondria. Caspase 9 is involved in the caspase activation cascade responsible for apoptosis execution and cleaves/activates Caspase 3 and Caspase 6. Caspase 9 is inhibited by the dominant negative isoform, BclXL, cIAP1, cIAP2, XIAP, and Livin. This caspase becomes activated when recruited to Apaf1/cytochrome c complex, and following cleavage by Apaf1, granzyme B, Caspase 3, possibly Caspase 8 and Caspase 10 into large p37 and small p10 subunits. Caspase 9 interacts with BIRC7 and has been shown to cleave PARP and vimentin.?

Gene ID:	842
UniProt:	P55211
Pathways:	MAPK Signaling , RTK Signaling , Apoptosis , Caspase Cascade in Apoptosis , Fc-epsilon Receptor Signaling Pathway , EGFR Signaling Pathway , Neurotrophin Signaling Pathway , Positive Regulation of Endopeptidase Activity

Application Details

Application Notes:	FCM 1:20-100
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months