

## Datasheet for ABIN6988676

## anti-STAT3 antibody (pSer727) (PE)



Go to Product page

_						
	V	$\triangle$	r۱	/1	$\triangle$	Λ/
	' V '		ΙV			v v

Quantity:	100 μL
Target:	STAT3
Binding Specificity:	pSer727
Reactivity:	Human, Rat, Rabbit
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STAT3 antibody is conjugated to PE
Application:	Western Blotting (WB), Flow Cytometry (FACS)
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human STAT3 around the
	phosphorylation site of Ser727
laatuna	
Isotype:	IgG
Cross-Reactivity:	lgG Human, Rabbit, Rat
Cross-Reactivity:	Human, Rabbit, Rat
Cross-Reactivity:  Predicted Reactivity:	Human, Rabbit, Rat  Mouse,Dog,Cow,Sheep,Pig,Chicken,Guinea Pig
Cross-Reactivity:  Predicted Reactivity:  Purification:	Human, Rabbit, Rat  Mouse,Dog,Cow,Sheep,Pig,Chicken,Guinea Pig
Cross-Reactivity:  Predicted Reactivity:  Purification:  Target Details	Human, Rabbit, Rat  Mouse,Dog,Cow,Sheep,Pig,Chicken,Guinea Pig  Purified by Protein A.

## **Target Details**

Larget Details		
Background:	Synonyms: APRF, HIES, ADMIO, Signal transducer and activator of transcription 3, Acute-phase response factor, STAT3  Background: Signal transducer and transcription activator that mediates cellular responses to interleukins, KITLG/SCF and other growth factors. May mediate cellular responses to activated FGFR1, FGFR2, FGFR3 and FGFR4. Binds to the interleukin-6 (IL-6)-responsive elements identified in the promoters of various acute-phase protein genes. Activated by IL31 through IL31RA. Cytoplasmic STAT3 represses macroautophagy by inhibiting EIF2AK2/PKR activity. Plays an important role in host defense in methicillin-resistant S.aureus lung infection by regulating the expression of the antimicrobial lectin REG3G (By similarity).	
Gene ID:	6774	
UniProt:	P40763	
Pathways:	JAK-STAT Signaling, RTK Signaling, Interferon-gamma Pathway, Neurotrophin Signaling Pathway, Dopaminergic Neurogenesis, Response to Growth Hormone Stimulus, Carbohydrate Homeostasis, Stem Cell Maintenance, Hepatitis C, Protein targeting to Nucleus, Feeding Behaviour, CXCR4-mediated Signaling Events, Signaling of Hepatocyte Growth Factor Receptor	
Application Details		
Application Notes:	FCM 1:20-100	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 a 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Ct - u- v-	00.00	

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

-20 °C

12 months

Storage:

Expiry Date:

Storage Comment: